

## University of Groningen

### **Association between administered oxygen, arterial partial oxygen pressure and mortality in mechanically ventilated intensive care unit patients**

de Jonge, Evert; Peelen, Linda; Keijzers, Peter J.; Joore, Hans; de Lange, Dylan; van der Voort, Peter H. J.; Bosman, Robert J.; de Waal, Ruud A. L.; Wesselink, Ronald; de Keizer, Nicolette F.

*Published in:*  
Critical Care

*DOI:*  
[10.1186/cc7150](https://doi.org/10.1186/cc7150)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2008

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

de Jonge, E., Peelen, L., Keijzers, P. J., Joore, H., de Lange, D., van der Voort, P. H. J., Bosman, R. J., de Waal, R. A. L., Wesselink, R., & de Keizer, N. F. (2008). Association between administered oxygen, arterial partial oxygen pressure and mortality in mechanically ventilated intensive care unit patients. *Critical Care*, 12(6), [R156]. <https://doi.org/10.1186/cc7150>

#### **Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### **Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Research

## Open Access

**Association between administered oxygen, arterial partial oxygen pressure and mortality in mechanically ventilated intensive care unit patients**Evert de Jonge<sup>1</sup>, Linda Peelen<sup>2,3</sup>, Peter J Keijzers<sup>4</sup>, Hans Joore<sup>4</sup>, Dylan de Lange<sup>4</sup>, Peter HJ van der Voort<sup>5</sup>, Robert J Bosman<sup>5</sup>, Ruud AL de Waal<sup>6</sup>, Ronald Wesselink<sup>7</sup> and Nicolette F de Keizer<sup>2</sup><sup>1</sup>Department of Intensive Care, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands<sup>2</sup>Department of Medical Informatics, Academic Medical Center, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands<sup>3</sup>Department of Epidemiology, Julius Center for Health Sciences and Primary Care, University Medical Center, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands<sup>4</sup>Intensive Care, University Medical Center, Heidelberglaan 100, 3584 CX, Utrecht, The Netherlands<sup>5</sup>Intensive Care, Onze Lieve Vrouwe Gasthuis, Oosterpark 91091 AC Amsterdam, The Netherlands<sup>6</sup>Intensive Care, Amphia Medical Center, Molengracht 21, 4818 CK Breda, The Netherlands<sup>7</sup>Intensive Care, St. Antonius Hospital Koekoekslaan 1, 3435 CM Nieuwegein, The NetherlandsCorresponding author: Evert de Jonge, [e.dejonge@amc.uva.nl](mailto:e.dejonge@amc.uva.nl)

Received: 4 Jul 2008 Revisions requested: 8 Sep 2008 Revisions received: 12 Nov 2008 Accepted: 10 Dec 2008 Published: 10 Dec 2008

*Critical Care* 2008, **12**:R156 (doi:10.1186/cc7150)This article is online at: <http://ccforum.com/content/12/6/R156>© 2008 de Jonge *et al.*; licensee BioMed Central Ltd.This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**Abstract****Introduction** The aim of this study was to investigate whether in-hospital mortality was associated with the administered fraction of oxygen in inspired air (FiO<sub>2</sub>) and achieved arterial partial pressure of oxygen (PaO<sub>2</sub>).**Methods** This was a retrospective, observational study on data from the first 24 h after admission from 36,307 consecutive patients admitted to 50 Dutch intensive care units (ICUs) and treated with mechanical ventilation. Oxygenation data from all admission days were analysed in a subset of 3,322 patients in 5 ICUs.**Results** Mean PaO<sub>2</sub> and FiO<sub>2</sub> in the first 24 h after ICU admission were 13.2 kPa (standard deviation (SD) 6.5) and 50% (SD 20%) respectively. Mean PaO<sub>2</sub> and FiO<sub>2</sub> from all admission days were 12.4 kPa (SD 5.5) and 53% (SD 18). Focusing on oxygenation in the first 24 h of admission, in-hospital mortality was shown to be linearly related to FiO<sub>2</sub> valueand had a U-shaped relationship with PaO<sub>2</sub> (both lower and higher PaO<sub>2</sub> values were associated with a higher mortality), independent of each other and of Simplified Acute Physiology Score (SAPS) II, age, admission type, reduced Glasgow Coma Scale (GCS) score, and individual ICU. Focusing on the entire ICU stay, in-hospital mortality was independently associated with mean FiO<sub>2</sub> during ICU stay and with the lower two quintiles of mean PaO<sub>2</sub> value during ICU stay.**Conclusions** Actually achieved PaO<sub>2</sub> values in ICU patients in The Netherlands are higher than generally recommended in the literature. High FiO<sub>2</sub>, and both low PaO<sub>2</sub> and high PaO<sub>2</sub> in the first 24 h after admission are independently associated with in-hospital mortality in ICU patients. Future research should study whether this association is causal or merely a reflection of differences in severity of illness insufficiently corrected for in the multivariate analysis.**Introduction**

It is generally acknowledged that mechanical ventilation may cause or exacerbate lung damage in critically ill patients with acute lung injury (ALI) or acute respiratory distress syndrome (ARDS). Many studies have examined the effects of different

settings of ventilation, such as low vs high tidal volumes, prone positioning and high-frequency oscillation on outcome of intensive care unit (ICU) patients [1]. Lung-protective mechanical ventilation strategies in patients with ALI/ARDS, applying lower tidal volumes and sufficient levels of positive end expira-

APACHE II: Acute Physiology and Chronic Health Evaluation II; FiO<sub>2</sub>: fraction of oxygen in the inspired air; MPM II: Mortality Prediction Model II; NICE: National Intensive Care Evaluation; PaO<sub>2</sub>: partial pressure of oxygen; SAPS II: Simplified Acute Physiology Score II; SOFA: Sequential Organ Failure Assessment.

tory pressure (PEEP) [2,3], have been shown to improve outcome.

The mode of mechanical ventilation and the oxygenation targets may influence the outcome for patients. Traditionally, arterial oxygen concentration (measured as partial oxygen pressure,  $\text{PaO}_2$ ) and oxygen saturation by pulse oximetry are used as targets. Common recommendations for oxygenation propose  $\text{PaO}_2$  values to be between 7.3 and 10.6 kPa [2,4]. The deleterious effects of hypoxia are well known and physicians may be mostly concerned about avoiding hypoxia and give additional oxygen 'to be on the safe side'. Hyperoxia, however, is also to be avoided as oxygen may be toxic. First, it is long known that high fraction of oxygen in inspired air ( $\text{FiO}_2$ ) may be toxic for the lungs. In animals, prolonged hyperoxia causes histopathological changes similar to those seen in ARDS [5]. Baboons exposed to 100% oxygen demonstrated a progressive reduction in forced vital capacity and functional residual capacity [6] and proliferative epithelial changes and interstitial fibrosis [7]. In healthy humans, exposure to 100% oxygen may lead to atelectasis, impaired mucociliary clearance and tracheobronchitis, alveolar protein leakage and enhanced expression of leukotrienes by alveolar macrophages and increases in alveolar neutrophils [8]. Apart from its effects on the lungs, oxygen may also lead to systemic toxicity. It has been associated with an increase in vascular resistance and a decrease in cardiac output [9]. Hyperoxia may result in the generation of central nervous system, hepatic and pulmonary free radicals. Cardiopulmonary resuscitation following cardiac arrest in a canine model is associated with a worsened neurologic outcome when performed in the presence of hyperoxia vs normoxia [8,10].

The aim of the present study was to describe the present oxygenation targets applied in ICUs in The Netherlands, and to determine whether outcome of ICU patients was associated with differences in administered oxygen ( $\text{FiO}_2$ ) or achieved arterial  $\text{PaO}_2$ .

## Materials and methods

### Patient data

This study is based on retrospective analysis of all consecutive patients admitted between 1 January 1999 and 30 June 2006 to the ICUs of 50 university, teaching and non-teaching hospitals in The Netherlands who were on mechanical ventilation within the first 24 h after ICU admission. Data were collected as part of the Dutch National Intensive Care Evaluation (NICE) registry. Within this registry data collection takes place in a standardised manner according to strict definitions and is subject to stringent data quality checks, which has been shown to result in a high quality of data [11]. The data have been encrypted in a way that all patient identifying information, such as name and patient identification number, has been removed. In The Netherlands, there is no need to obtain consent to make use of registries when patient identifying information is not

used. According to the Dutch Medical Research Involving Human Subjects Act, there is no need for approval by ethical committees [12]. The NICE initiative is officially registered according to the Dutch Personal Data Protection Act.

The variables are used to calculate probabilities of death for each patient using the Acute Physiology and Chronic Health Evaluation (APACHE) II [13], the Simplified Acute Physiology Score (SAPS) II [14], and the Mortality Probability Models (MPM) II at admission and 24-h scoring systems [15]. In this study the SAPS II was used for case mix adjustment as previous research has shown that this scoring system fits best to the patient population of the NICE registry [16]. The database contains 108 demographic, diagnostic and physiologic variables collected within the first 24 h of ICU admission and outcome data on ICU and in-hospital mortality.

In analogy with the exclusion criteria commonly used in analyses based on the SAPS II scoring system, patients admitted after cardiac surgery, patients admitted with severe burns and patients aged under 18 were excluded from the analyses. For patients with multiple ICU admissions during a hospitalisation period only the first ICU admission was used.

In the analyses focusing on oxygenation in the first 24 h of ICU stay, information of all patients was used. For the analyses related to oxygenation during the entire ICU stay a selection of the patients was used, as only five of the ICUs participating in the NICE registry provide information to the registry database on the patient's condition on a daily basis using the Sequential Organ Failure Assessment (SOFA) score [17]. For this analysis only patients with a minimum length of ICU stay of 3 days were included into the analyses. Mean  $\text{PaO}_2$  and mean  $\text{FiO}_2$  values were calculated based on the entire ICU stay.

If more than one blood gas analysis was available for a patient during the first 24 h after ICU admission,  $\text{PaO}_2$ ,  $\text{FiO}_2$  and partial  $\text{CO}_2$  ( $\text{PaCO}_2$ ) values were from the arterial sample with the lowest  $\text{PaO}_2/\text{FiO}_2$  ratio. Likewise, oxygenation data in the SOFA scores was based on samples with lowest  $\text{PaO}_2/\text{FiO}_2$  ratios in the particular 24 h period.

### Statistical analyses

The relation between the oxygenation parameters and in-hospital mortality was assessed using logistic regression analysis.

As  $\text{PaO}_2$  and  $\text{FiO}_2$  are both continuous variables, univariate regression analyses using polynomial functions and spline functions [18,19] were performed to investigate the relation between each of these variables and in-hospital mortality. For  $\text{PaO}_2$  a model including  $\text{PaO}_2$  as a natural spline with five degrees of freedom turned out to result in the best fit. To enhance interpretation of the results, in subsequent analyses  $\text{PaO}_2$  was categorised. As no standard cut-off points are in use for  $\text{PaO}_2$ , categorisation of  $\text{PaO}_2$  values into five catego-

ries was based on the distribution of the data, using quintiles as cut-off values between categories. For  $\text{FiO}_2$  inclusion into the model in a linear fashion showed to be optimal fit.

Multivariable logistic regression analyses were performed both for the first 24 h of ICU stay and for the entire ICU stay. Potential confounders in the association of oxygenation with hospital mortality (age, SAPS II, Glasgow Coma Scale (GCS) score below 15 and admission type) were included into the models. Also a specific variable denoting the 'hospital' was included into these models to correct for potential differences in overall in-hospital mortality between the five hospitals. In the model focusing on the entire ICU stay, the  $\text{PaO}_2/\text{FiO}_2$  ratio during the first 24 h of admission was added as additional confounder. In the modelling process the presence of multicollinearity between the oxygenation parameters was verified based on the standard errors of the parameters in the model.

The Standardised Mortality Ratio (SMR) was calculated as the ratio of the number of observed deaths to the number of deaths expected according to the SAPS II model [13].

In all analyses a p value of 0.05 was considered to represent a statistically significant difference. The analyses were performed using SPSS version 14.0 (SPSS Inc., Chicago, IL, USA) and S-plus version 7.0 (Insightful Corp., Seattle, WA, USA).

## Results

### Analysis of data from first 24 h after admission

In total, 36,307 patients from 50 ICUs were included in the analysis. All patients were treated with mechanical ventilation within the first 24 h after ICU admission. Data on the severity of illness, reason for admission and referring specialty are given in Table 1. Mean  $\text{PaO}_2$  was 13.2 kPa (standard deviation (SD) 6.5). Mean  $\text{FiO}_2$  was 50% (SD 20). Regression analysis using  $\text{PaO}_2$  as a continuous variable showed that with increasing  $\text{PaO}_2$  the average in-hospital mortality first decreased and subsequently started to rise again (Figure 1). Figure 2 denotes the relation between  $\text{PaO}_2$  and in-hospital mortality when correcting for SAPS II (by means of the SMRs). Figures 3 and 4 show the association between SMR and  $\text{FiO}_2$  or  $\text{PaO}_2/\text{FiO}_2$  ratio respectively. Multivariate regression analysis indicated that the U-shaped relation between  $\text{PaO}_2$  and mortality (modelled using a spline function) remained significant after correction for age, admission type, GCS score and severity of illness measured with the SAPS II. Multicollinearity was not found to be present for  $\text{FiO}_2$  and  $\text{PaO}_2$  values. Table 2 presents the odds ratios for  $\text{FiO}_2$  and the  $\text{PaO}_2$  quintiles in a multivariate regression model, indicating that in-hospital mortality was associated with  $\text{FiO}_2$  and  $\text{PaO}_2$  values.

### Analysis of data from all ICU admission days

For the analysis regarding entire ICU stay 3,322 patients from 5 ICUs were included. Characteristics of these patients are

given in Table 1. Mean  $\text{PaO}_2$  during ICU stay was 12.4 kPa (SD 5.5). Mean  $\text{FiO}_2$  during ICU stay was 53% (SD 18). The results of the multivariate analysis are shown in Table 3. Mean  $\text{FiO}_2$  value and a mean  $\text{PaO}_2$  value lower than 10.6 kPa were associated with a higher mortality. This association was independent of the potential confounders age, SAPS II, abnormal GCS score,  $\text{PaO}_2/\text{FiO}_2$  ratio at admission, admission type and hospital.

## Discussion

We found that administration of high  $\text{FiO}_2$  values in ICU patients was associated with increased in-hospital mortality. This association was found for  $\text{FiO}_2$  values in the first 24 h after admission and also for mean  $\text{FiO}_2$  during all admission days. The increased risk in patients with high  $\text{FiO}_2$  remained after correcting for SAPS II, admission type, reduced GCS score and pulmonary dysfunction measured as  $\text{PaO}_2/\text{FiO}_2$  ratio. This suggests that the administration of oxygen itself could be deleterious, and that the association between high  $\text{FiO}_2$  and mortality cannot be explained by the confounding issue that highest  $\text{FiO}_2$  levels are administered in patients with severe pulmonary dysfunction.

Our observations are in accordance with prior experimental studies showing the potential toxicity of high fractions of inspired oxygen [5]. Administration of supplemental oxygen can cause lung damage. This risk is especially high in prematurely born infants, probably attributable to inadequate host defences, underdeveloped lungs and immature antioxidant systems [20]. Exposure to hyperoxia leads to diffuse pulmonary damage characterised by an extensive inflammatory response and destruction of the alveolar-capillary barrier leading to oedema, impaired gas exchange and respiratory failure [21]. Mouse lungs exposed to > 90% oxygen for 48 h were more susceptible to ventilator-induced lung injury than those exposed to room air [22]. Hyperoxia also aggravates pulmonary injury following artificial ventilation in rats using high tidal volumes [23]. Furthermore, hyperoxia impairs the innate immune response by decreased macrophage function, impaired bacterial killing and increased susceptibility to pneumonia in a *Klebsiella pneumoniae* model [24]. Lung injury is likely to be initiated when the rates of generation of reactive oxygen species (ROS) are increased beyond the capacities of the antioxidant defences, such as the enzymes glutathione, superoxide dismutase and catalase. Mitochondrial mediated cell injury by ROS has been identified as a critical event in both apoptotic and necrotic forms of cell death in hyperoxia [25]. Another organ that may be injured by hyperoxia is the kidney. Hyperoxic reperfusion exacerbates renal dysfunction and histopathologic injury after 30 min of complete normothermic ischaemia in rabbits. This hyperoxia associated dysfunction was prevented by the administration of the radical scavenger allopurinol [26], suggesting that oxidative injury by ROS plays a role in post-ischaemic renal failure.

**Table 1****Characteristics of patients**

	Analysis of data from first 24 h after admission	Analysis of data from all admission days
No of patients	36,307	3,322
Male (%)	60.1	60.3
Age in years <sup>a</sup>	62.5 ± 16.1	62.4 ± 15.9
SAPS II <sup>a</sup>	42.7 ± 18.4	47.6 ± 15.7
SAPS II predicted mortality <sup>a</sup>	0.34 ± 0.28	0.42 ± 0.27
GCS score below 15 (%)	33.1	33.6
PaO <sub>2</sub> at admission (kPa) <sup>a</sup>	13.2 ± 6.5 <sup>b</sup>	12.5 ± 5.5 <sup>c</sup>
FiO <sub>2</sub> (%) <sup>a</sup>	50.4 ± 19.9 <sup>b</sup>	53.1 ± 18.7 <sup>c</sup>
PaO <sub>2</sub> /FiO <sub>2</sub> ratio (kPa) <sup>a</sup>	29.1 ± 15.0 <sup>b</sup>	24.6 ± 12.6 <sup>c</sup>
Admission type (%):		
Medical	48.4	61.0
Unplanned surgery	22.6	23.0
Planned surgery	28.9	16.0
Referring specialty (%):		
Internal medicine	16.4	16.7
Cardiology	10.8	17.9
Pulmonary disease	7.4	10.4
Neurology	5.7	6.4
Surgery	33.2	30.6
Cardiothoracic surgery	6.4	1.2
Neurosurgery	6.9	6.1
Other	13.1	10.7
ICU mortality	23.0	21.0
In-hospital mortality (%)	31.1	32.7

<sup>a</sup>Mean ± standard deviation (SD); <sup>b</sup>PaO<sub>2</sub>, FiO<sub>2</sub> and PaO<sub>2</sub>/FiO<sub>2</sub> ratio from sample with lowest PaO<sub>2</sub>/FiO<sub>2</sub> ratio within 24 h after admission; <sup>c</sup>mean value of all admission days. Per day PaO<sub>2</sub>, FiO<sub>2</sub> and PaO<sub>2</sub>/FiO<sub>2</sub> ratio was taken from sample with lowest PaO<sub>2</sub>/FiO<sub>2</sub> ratio.

GCS, Glasgow Coma Scale; FiO<sub>2</sub>, fraction of oxygen in inspired air; ICU, intensive care unit; PaO<sub>2</sub>, partial oxygen pressure; SAPS II, Simplified Acute Physiology Score II.

Several studies focused on the role of high reperfusion oxygen tensions following cardiac arrest and resuscitation. In a canine model of 10 min of cardiac arrest, resuscitation with 21% vs 100% inspired O<sub>2</sub> resulted in lower levels of oxidised brain lipids and improved neurological outcome [27]. In another study using the same canine model, it was shown that resuscitation with 100% O<sub>2</sub> resulted in impaired hippocampal neuronal metabolism [28]. Proposed pathogenetic mechanisms of hyperoxia induced reperfusion injury of the brain include increased production of ROS, a high ratio of oxidised over reduced glutathione [29] and increased nitric oxide production by endothelium and neuron derived nitric oxide synthase [30].

Many studies investigated the use of 100% vs 21% oxygen for resuscitation in depressed newborn infants (that is, infants with apnoea or relative bradycardia at birth). A systematic review and meta-analysis of 10 studies reported a significant reduction in the risk of neonatal mortality and a trend towards a reduction in severe encephalopathy in newborns resuscitated with 21% O<sub>2</sub>. The reduction in mortality was also found in a subgroup analysis only including strictly randomised controlled trials and in a subgroup of studies enrolled in European countries with a lower risk of mortality than in less developed countries [31].

Human clinical studies evaluating the effects of hyperoxia in critically ill adult patients are lacking. The effects of hyperoxia in non-ICU settings are not clear. A reduction in surgical site

**Table 2**

**Adjusted odds ratios for partial oxygen pressure (PaO<sub>2</sub>) and fraction of oxygen in inspired air (FiO<sub>2</sub>) resulting from a multivariate regression analysis on data from the first 24 h after ICU admission**

Covariate	Odds ratio	95% Confidence interval
PaO <sub>2</sub> in kPa:		
< 8.9 (n = 6,937)	1.12	1.03 to 1.21
8.9 to 10.6 (reference category) (n = 7,466)	1	
10.6 to 12.6 (n = 6,430)	1.11	1.02 to 1.21
12.6 to 16.4 (n = 7,278)	1.08	1.00 to 1.18
≥ 16.4 (n = 8,196)	1.23	1.13 to 1.34
FiO <sub>2</sub> (per 10%)	1.12	1.10 to 1.13

Odds ratio after adjustment for the following potential confounders: age, SAPS II, GCS score below 15, admission type, individual hospital. The equation of the model is:  $\text{Logit}(p) = -5.419 + 0.059 \times \text{age (per 5 years)} + 0.066 \times \text{SAPS II} + 0.070 \times I(\text{GCS} < 15) + 0.221 \times I(\text{admission type} = \text{urgent}) + 0.453 \times I(\text{admission type} = \text{medical}) + \beta_{\text{hosp}} + 0.105 \times \text{FiO}_2 \text{ (per 10\%)} + 0.109 \times I(\text{PaO}_2 < 67) + 0.109 \times I(80 \leq \text{PaO}_2 < 95) + 0.079 \times I(95 \leq \text{PaO}_2 < 123) + 0.206 \times I(\text{PaO}_2 \geq 123)$  Probability of in-hospital death =  $e^{(\text{logit})} / (1 + e^{(\text{logit})})$ . Median  $\beta_{\text{hosp}}$  for individual hospitals was -0.12 (IQR -0.43 to 0.05).


GCS, Glasgow Coma Scale; ICU, intensive care unit; IQR, interquartile range; SAPS II, Simplified Acute Physiology Score II.

infections by the use of hyperoxia has been reported by one study group [32], while others reported more surgical site infections in patients treated with hyperoxia [33].


An alternative explanation for the association between oxygenation and mortality in ICU patients could be that common criteria for weaning from mechanical ventilation are based on FiO<sub>2</sub> and PEEP levels. High FiO<sub>2</sub> and PEEP, both leading to high PaO<sub>2</sub> values, may delay weaning from mechanical ventilation, thus negatively influencing outcome in ICU patients. Also, we cannot exclude that high PaO<sub>2</sub> values were achieved by

more invasive ventilation strategies, potentially being more injurious to the patients.


Interestingly, apart from FiO<sub>2</sub> values, there was also a U-shaped association between achieved arterial oxygen tension (PaO<sub>2</sub>) during the first 24 h after ICU admission and mortality with higher mortality in patients with either a very low or high PaO<sub>2</sub>. That mortality is higher in patients with very low PaO<sub>2</sub> is not unexpected and possibly related to ischaemia or to selection of the sickest patients. However, mortality was also higher in patients with highest PaO<sub>2</sub> values, suggesting the possibility of systemic oxygen toxicity.

**Figure 1**

**In-hospital mortality by partial oxygen pressure (PaO<sub>2</sub>) (kPa).** Values were taken from blood gas analysis with lowest PaO<sub>2</sub>/fraction of oxygen in inspired air (FiO<sub>2</sub>) ratio in the first 24 h after intensive care unit (ICU) admission. The sizes of the circles represent the number of patients with the same PaO<sub>2</sub> value. The curve represents the predicted mortality using the logistic regression equation in which the PaO<sub>2</sub> value was incorporated using a spline function.


**Figure 2**

**Standardised mortality ratio (SMR) by partial oxygen pressure (PaO<sub>2</sub>) (kPa).** PaO<sub>2</sub> values were taken from blood gas analysis with lowest PaO<sub>2</sub>/fraction of oxygen in inspired air (FiO<sub>2</sub>) ratio in the first 24 h after intensive care unit (ICU) admission. PaO<sub>2</sub> values are categorised as quintiles. Error bars represent 95% confidence intervals.

**Figure 3**

**Standardised mortality ratio (SMR) by fraction of oxygen in inspired air (FiO<sub>2</sub>).** FiO<sub>2</sub> values were taken from blood gas analysis with lowest partial oxygen pressure (PaO<sub>2</sub>)/FiO<sub>2</sub> ratio in the first 24 h after intensive care unit (ICU) admission. FiO<sub>2</sub> values are categorised as quintiles. Error bars represent 95% confidence intervals.

In our analysis of mean oxygenation during all admission days, we again found a linear association between mortality and FiO<sub>2</sub> values. Low PaO<sub>2</sub>s were also associated with higher mortality but high PaO<sub>2</sub>s were not. The shape of the association between PaO<sub>2</sub> and mortality was hard to assess. In our data a linear association appeared to best fit the data (data not shown). The number of patients included in this analysis was only 3,322. Only 2% of the patients had a mean PaO<sub>2</sub> higher than 20.0 kPa. Thus, the power of our study may have been too low to detect an association between high mean PaO<sub>2</sub> values during the ICU stay and increased mortality.

**Figure 4**

**Standardised mortality ratio (SMR) by lowest partial oxygen pressure (PaO<sub>2</sub>)/fraction of oxygen in inspired air (FiO<sub>2</sub>) ratio (kPa) in the first 24 h after intensive care unit (ICU) admission.** PaO<sub>2</sub>/FiO<sub>2</sub> ratio values are categorised as quintiles. Error bars represent 95% confidence intervals.

There are limitations to this study. Most importantly, it was a retrospective observational study and the association between mortality and oxygenation is not necessarily causal. Although the association appeared to be independent of a number of potential confounding covariates, we cannot exclude that, despite our efforts, there are still differences in case mix associated with oxygenation that are not taken into account in our multivariate analyses. It is possible that physicians recognised some marker of severity that was not represented in our attempts to adjust for severity, and that they purposefully gave higher concentrations of oxygen to achieve higher levels of PaO<sub>2</sub> in these high-risk patients.

**Table 3**

**Adjusted odds ratios for mean partial oxygen pressure (PaO<sub>2</sub>) value and mean fraction of oxygen in inspired air (FiO<sub>2</sub>) during ICU stay resulting from a multivariate regression analysis on data from the entire ICU stay**

Covariate	Odds ratio	95% Confidence interval
Mean PaO <sub>2</sub> in kPa:		
< 8.9 (n = 402)	1.63	1.16 to 2.3
8.9 to 10.6 (n = 871)	1.51	1.18 to 1.96
10.6 to 12.6 (n = 970)	1.25	0.99 to 1.57
12.6 to 16.4 (reference category) (n = 841)	1	
> 16.4 (n = 238)	1.04	0.64 to 1.68
Mean FiO <sub>2</sub> (per 10%)	1.63	1.47 to 1.81

Odds ratio after adjustment for the following potential confounders: age, SAPS II, GCS score below 15, admission type, PaO<sub>2</sub>/FiO<sub>2</sub> ratio at admission, and hospital. The equation of the eventual model is:  $\text{Logit}(p) = -7.060 + 0.090 \times \text{age (per 5 years)} + 0.049 \times \text{SAPS II} + 0.054 \times \text{I}(\text{GCS} < 15) + 0.015 \times \text{I}(\text{admission type} = \text{urgent}) + 0.161 \times \text{I}(\text{admission type} = \text{medical}) + 0.004 \times \text{PaO}_2/\text{FiO}_2 - 1.114 \times \text{I}(\text{hospital} = 2) - 0.060 \times \text{I}(\text{hospital} = 3) - 0.285 \times \text{I}(\text{hospital} = 4) - 0.618 \times \text{I}(\text{hospital} = 5) + 0.488 \times \text{mean FiO}_2 \text{ (per 10\%)} + 0.492 \times \text{I}(\text{mean PaO}_2 < 67) + 0.417 \times \text{I}(67 \leq \text{PaO}_2 < 80) + 0.221 \times \text{I}(80 \leq \text{PaO}_2 < 95) + 0.038 \times \text{I}(\text{PaO}_2 \geq 123)$ . Probability of in-hospital death =  $e^{(\text{logit})}/(1 + e^{(\text{logit})})$ . GCS, Glasgow Coma Scale; ICU, intensive care unit; SAPS II, Simplified Acute Physiology Score II.

The three potential confounders that we corrected for (age, reduced GCS score, and admission type) are part of the SAPS II that was also included as covariate in the multivariate analysis. We have repeated the analyses without these three variables, adjusting for SAPS II only. This yielded similar results for the association between in-hospital mortality and PaO<sub>2</sub> and FiO<sub>2</sub> respectively.

We corrected for pulmonary dysfunction by including PaO<sub>2</sub>/FiO<sub>2</sub> ratio at admission in the multivariate analysis of data from all admission days. PaO<sub>2</sub>/FiO<sub>2</sub> ratio was not included in the analysis of data from the first 24 h after ICU admission, because including PaO<sub>2</sub>, FiO<sub>2</sub> and PaO<sub>2</sub>/FiO<sub>2</sub> ratio, all from the same arterial blood sample, would introduce problems by colinearity of the data. In this population, however, we performed a separate multivariate analysis substituting PaO<sub>2</sub>/FiO<sub>2</sub> ratio for PaO<sub>2</sub> values. Again, FiO<sub>2</sub> appeared to be a predictor of mortality, also independent of PaO<sub>2</sub>/FiO<sub>2</sub> ratio (OR 1.15, 95% CI 1.14 to 1.17, model not shown). PaO<sub>2</sub>/FiO<sub>2</sub> ratio is not only influenced by pulmonary dysfunction, but also by ventilator settings, such as PEEP levels. As PEEP was not part of the NICE data collection, we could not include this possible confounder in our analysis. Prospective, controlled trials are necessary to show a causal relationship between high FiO<sub>2</sub>s and mortality.

As the association between PaO<sub>2</sub> and mortality was U-shaped, we categorised PaO<sub>2</sub> values for the multivariate analysis using quintiles as categories (as no standard categorisation is available). The boundaries of quintiles are chosen arbitrarily and may not be the optimal cut-off levels to discriminate between patients with low and high risk of mortality. Therefore, we repeated the same multivariate analysis (model not shown) on data from the first 24 h after ICU admission using PaO<sub>2</sub> values categorised as deciles and found similar results.

Another finding from our study is the fact that in most patients the achieved PaO<sub>2</sub> values are higher than the targets commonly recommended [2,4]. Although oxygen toxicity is a well known entity [34], FiO<sub>2</sub>s up to 0.5 are commonly considered 'safe' by physicians [5]. It appears that physicians are more concerned about avoiding hypoxia and ischaemia than about the risks of hyperoxia. In The Netherlands, no formal guidelines for oxygenation targets are available. This may be related to the fact that the influence of oxygenation targets has never been studied making it impossible to provide evidence-based recommendations. Based on other observational studies, it may well be that also in other countries actual PaO<sub>2</sub>s in ICU patients are higher than recommended [35,36]

## Conclusion

High fractions of oxygen in the inspired air and high PaO<sub>2</sub> values are associated with increased mortality in ICU patients. Actually achieved PaO<sub>2</sub> values in Dutch ICU patients are

higher than the PaO<sub>2</sub> targets in some recent international recommendations. Prospective interventional studies are necessary to find out whether the association between outcome and oxygenation is causal and to provide evidence-based guidelines on oxygenation targets.

## Key messages

- The weaning rate of catecholamines is usually chosen empirically by intensivists.
- Actually achieved PaO<sub>2</sub> values in Dutch ICU patients are higher than the PaO<sub>2</sub> targets given in recent international recommendations.
- High fractions of oxygen in the inspired air are associated with increased mortality in ICU patients on mechanical ventilation.
- Both low and high PaO<sub>2</sub> values in the first 24 hours after ICU admission were associated with increased mortality.
- Future interventional studies are required to find out whether these associations between oxygenation and outcome are causal or due to other confounding issues.

## Competing interests

During the period from 2002 to 2004 LP received an unrestricted educational grant from Eli Lilly Netherlands B.V. The study described in this manuscript was not conducted under the grant, and Eli Lilly Netherlands B.V. has not been involved in any part of the present study. All other authors declare that they have no competing interests.

## Authors' contributions

EdJ designed the study and drafted the manuscript. LP and NdK were involved in the set-up of the study, performed the statistical analyses and helped in interpreting the results and writing the manuscript. PK was involved in the set-up of the study, interpreting the results and writing the manuscript. JJ, DdL, PvdV, RB, RdW and RW were involved in interpreting the results and writing the manuscript. All authors read and approved the final manuscript.

## References

1. Fan E, Needham DM, Stewart TE: **Ventilatory management of acute lung injury and acute respiratory distress syndrome.** *JAMA* 2005, **294**:2889-2896.
2. The Acute Respiratory Distress Syndrome Network: **Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome.** *N Engl J Med* 2000, **342**:1301-1308.
3. Amato MB, Barbas CS, Medeiros DM, Magaldi RB, Schettino GP, Lorenzi-Filho G, Kairalla RA, Deheinzelin D, Munoz C, Oliveira R, Takagaki TY, Carvalho CR: **Effect of a protective-ventilation strategy on mortality in the acute respiratory distress syndrome.** *N Engl J Med* 1998, **338**:347-354.
4. Wheeler AP, Bernard GR: **Acute lung injury and the acute respiratory distress syndrome: a clinical review.** *Lancet* 2007, **369**:1553-1564.



5. Altemeier WA, Sinclair SE: **Hyperoxia in the intensive care unit: why more is not always better.** *Curr Opin Crit Care* 2007, **13**:73-78.
6. Fracica PJ, Knapp MJ, Piantadosi CA, Takeda K, Fulkerson WJ, Coleman RE, Wolfe WG, Crapo JD: **Responses of baboons to prolonged hyperoxia: physiology and qualitative pathology.** *J Appl Physiol* 1991, **71**:2352-2362.
7. Crapo JD, Hayatdavoudi G, Knapp MJ, Fracica PJ, Wolfe WG, Piantadosi CA: **Progressive alveolar septal injury in primates exposed to 60% oxygen for 14 days.** *Am J Physiol* 1994, **267**:L797-L806.
8. Kavanagh BP: **Goals and concerns for oxygenation in acute respiratory distress syndrome.** *Curr Opin Crit Care* 1998, **4**:16-20.
9. Lodato RF: **Decreased O<sub>2</sub> consumption and cardiac output during normobaric hyperoxia in conscious dogs.** *J Appl Physiol* 1989, **67**:1551-1559.
10. Zwemer CF, Whitesall SE, D'Alecy LG: **Hypoxic cardiopulmonary-cerebral resuscitation fails to improve neurological outcome following cardiac arrest in dogs.** *Resuscitation* 1995, **29**:225-236.
11. Arts D, de Keizer NF, Scheffer GJ, de Jonge E: **Quality of data collected for severity of illness scores in the Dutch National Intensive Care Evaluation (NICE) registry.** *Intensive Care Med* 2002, **28**:656-659.
12. The Central Committee on Research Involving Human Subjects (CCMO) [<http://www.ccmo-online.nl>]
13. Knaus WA, Draper EA, Wagner DP, Zimmerman JE: **APACHE II: a severity of disease classification system.** *Crit Care Med* 1985, **13**:818-829.
14. Le Gall JR, Lemeshow S, Saulnier F: **A new Simplified Acute Physiology Score (SAPS II) based on a European/North American multicenter study.** *JAMA* 1993, **270**:2957-2963.
15. Lemeshow S, Teres D, Klar J, Avrunin JS, Gehlbach SH, Rapoport J: **Mortality Probability Models (MPM II) based on an international cohort of intensive care unit patients.** *JAMA* 1993, **270**:2478-2486.
16. Peek N, Arts DG, Bosman RJ, Voort P van der, de Keizer NF: **External validation of prognostic models for critically ill patients required substantial sample sizes.** *J Clin Epidemiol* 2007, **60**:491-501.
17. Vincent JL, de MA, Cantraine F, Moreno R, Takala J, Suter PM, Sprung CL, Colardyn F, Blecher S: **Use of the SOFA score to assess the incidence of organ dysfunction/failure in intensive care units: results of a multicenter, prospective study. Working group on "sepsis-related problems" of the European Society of Intensive Care Medicine.** *Crit Care Med* 1998, **26**:1793-1800.
18. Smith PL: **Splines as a useful and convenient statistical tool.** *American Statistician* 1979, **33**:57-62.
19. Harrell FE: *Regression Modeling Strategies* New York, NY: Springer; 2001.
20. O'Donovan DJ, Fernandes CJ: **Mitochondrial glutathione and oxidative stress: implications for pulmonary oxygen toxicity in premature infants.** *Mol Genet Metab* 2000, **71**:352-358.
21. Crapo JD, Barry BE, Foscue HA, Shelburne J: **Structural and biochemical changes in rat lungs occurring during exposures to lethal and adaptive doses of oxygen.** *Am Rev Respir Dis* 1980, **122**:123-143.
22. Bailey TC, Martin EL, Zhao L, Veldhuizen RA: **High oxygen concentrations predispose mouse lungs to the deleterious effects of high stretch ventilation.** *J Appl Physiol* 2003, **94**:975-982.
23. Quinn DA, Moufarrej RK, Volokhov A, Hales CA: **Interactions of lung stretch, hyperoxia, and MIP-2 production in ventilator-induced lung injury.** *J Appl Physiol* 2002, **93**:517-525.
24. Baleeiro CE, Wilcoxon SE, Morris SB, Standiford TJ, Paine R III: **Sublethal hyperoxia impairs pulmonary innate immunity.** *J Immunol* 2003, **171**:955-963.
25. Wallace KB, Eells JT, Madeira VM, Cortopassi G, Jones DP: **Mitochondria-mediated cell injury. Symposium overview.** *Fundam Appl Toxicol* 1997, **38**:23-37.
26. Zwemer CF, Shoemaker JL Jr, Hazard SW III, Davis RE, Bartoletti AG, Phillips CL: **Hyperoxic reperfusion exacerbates postischemic renal dysfunction.** *Surgery* 2000, **128**:815-821.
27. Liu Y, Rosenthal RE, Haywood Y, Miljkovic-Lolic M, Vanderhoek JY, Fiskum G: **Normoxic ventilation after cardiac arrest reduces oxidation of brain lipids and improves neurological outcome.** *Stroke* 1998, **29**:1679-1686.
28. Richards EM, Fiskum G, Rosenthal RE, Hopkins I, McKenna MC: **Hyperoxic reperfusion after global ischemia decreases hippocampal energy metabolism.** *Stroke* 2007, **38**:1578-1584.
29. Gelfand SL, Vento M, Sastre J, Lust WD, Smith MA, Perry G, Walsh M, Martin R: **A new model of oxidative stress in rat pups.** *Neonatology* 2008, **94**:293-299.
30. Allen BW, Demchenko IT, Piantadosi CA: **Two faces of nitric oxide: implications for cellular mechanisms of oxygen toxicity.** *J Appl Physiol* 2008 in press.
31. Saugstad OD, Ramji S, Soll RF, Vento M: **Resuscitation of newborn infants with 21% or 100% oxygen: an updated systematic review and meta-analysis.** *Neonatology* 2008, **94**:176-182.
32. Greif R, Akca O, Horn EP, Kurz A, Sessler DI: **Supplemental perioperative oxygen to reduce the incidence of surgical-wound infection. Outcomes Research Group.** *N Engl J Med* 2000, **342**:161-167.
33. Pryor KO, Fahey TJ III, Lien CA, Goldstein PA: **Surgical site infection and the routine use of perioperative hyperoxia in a general surgical population: a randomized controlled trial.** *JAMA* 2004, **291**:79-87.
34. Mao C, Wong DT, Slutsky AS, Kavanagh BP: **A quantitative assessment of how Canadian intensivists believe they utilize oxygen in the intensive care unit.** *Crit Care Med* 1999, **27**:2806-2811.
35. Young MP, Manning HL, Wilson DL, Mette SA, Riker RR, Leiter JC, Liu SK, Bates JT, Parsons PE: **Ventilation of patients with acute lung injury and acute respiratory distress syndrome: has new evidence changed clinical practice?** *Crit Care Med* 2004, **32**:1260-1265.
36. Kalhan R, Mikkelsen M, Dedhiya P, Christie J, Gaughan C, Lanken PN, Finkel B, Gallop R, Fuchs BD: **Underuse of lung protective ventilation: analysis of potential factors to explain physician behavior.** *Crit Care Med* 2006, **34**:300-306.